



Sets non-thin at ∞ in \mathbb{C}^m

Tuyen Trung Truong

Department of Mathematics, Indiana University Bloomington, IN 47405, USA

ARTICLE INFO

Article history:

Received 10 December 2008

Available online 11 March 2009

Submitted by T. Ransford

Keywords:

Growth of entire functions

Non-thin at infinity

Pluricomplex Green function

Robin constant

Siciak extremal function

ABSTRACT

In this paper we define the notion of non-thin at ∞ as follows: Let E be a subset of \mathbb{C}^m . For any $R > 0$ define $E_R = E \cap \{z \in \mathbb{C}^m : |z| \leq R\}$. We say that E is non-thin at ∞ if

$$\lim_{R \rightarrow \infty} V_{E_R}(z) = 0$$

for all $z \in \mathbb{C}^m$, where V_E is the pluricomplex Green function of E . This definition of non-thinness at ∞ has good properties: If $E \subset \mathbb{C}^m$ is non-thin at ∞ and A is pluripolar then $E \setminus A$ is non-thin at ∞ ; if $E \subset \mathbb{C}^m$ and $F \subset \mathbb{C}^n$ are arbitrary sets, then E and F are non-thin at ∞ iff $E \times F \subset \mathbb{C}^m \times \mathbb{C}^n$ is non-thin at ∞ (see Lemma 2). The results of this paper extend some results in [J. Muller, A. Yavrian, On polynomials sequences with restricted growth near infinity, Bull. London Math. Soc. 34 (2002) 189–199] and [Dang Duc Trong, Tuyen Trung Truong, The growth at infinity of a sequence of entire functions of bounded orders, Complex Var. Elliptic Equ. 53 (8) (2008) 717–743].

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

Fix $m \in \mathbb{N}$, let \mathbb{C}^m be the usual m -dimensional complex Euclidean space. Before going into the main points, we recall some facts about the potential theory in \mathbb{C}^m . Let U be an open subset of \mathbb{C}^m . A function $u : U \rightarrow [-\infty, \infty)$ is called PSH in U (written $u \in PSH(U)$) if u is upper-semicontinuous and when restricted to any complex line $L \simeq \mathbb{C}$ then u is subharmonic (see [4]).

A $PSH(\mathbb{C}^m)$ function u is said to be of minimal growth if

$$u(z) - \log |z| \leq O(1), \quad \text{as } |z| \rightarrow \infty,$$

here $|z|$ is the usual Euclidean norm of an element $z \in \mathbb{C}^m$. We denote the set of all such functions by \mathbb{L} .

Let E be a subset in \mathbb{C}^m . Then the pluricomplex Green function of the set E with pole at infinity (see [4, p. 184]) is

$$V_E(z) = \sup\{u(z) : u \in \mathbb{L}, u|_E \leq 0\} \quad (z \in \mathbb{C}^m).$$

V_E is also called the Siciak extremal function of the set E . By definition we see that $V_E \geq 0$.

For any function $u : \mathbb{C}^m \rightarrow [-\infty, \infty)$ we define its upper-semicontinuous regularization u^* (see [4]) by

$$u^*(z) = \limsup_{\zeta \rightarrow z} u(\zeta).$$

Let $V_E^*(z)$ be the regularization of the Green function $V_E(z)$. It is well known (see [4]) that there are only three cases:

E-mail address: truongt@indiana.edu.

Case 1: $V_E^*(z) \equiv +\infty$. Then E is pluripolar.

Case 2: $V_E^*(z) \in \mathbb{L}$, and $V_E^* \not\equiv 0$.

Case 3: $V_E^*(z) \equiv 0$.

When $m = 1$ then $V_E^*(z) \equiv 0$ iff the set E is non-thin at ∞ , or equivalently the set $E^* = \{z: 1/z \in E\}$ is non-thin at 0 (see [7,2]). Recall that a subset E of \mathbb{C}^m is pluri-thin (or thin for brevity) at a point $a \in \mathbb{C}^m$ if either a is not a limit point of E or there is a neighborhood U of a and a function $u \in PSH(U)$ such that

$$\limsup_{z \rightarrow a, z \in E \setminus a} u(z) < u(a).$$

If E is not thin then it is called non-thin.

In [7] the authors proved the following result (see also [2, p. 270]).

Proposition 1. *Let E be a closed subset of \mathbb{C} . Then the following four statements are equivalent:*

1. E is non-thin at ∞ .
2. For any $z \in \mathbb{C}$ we have

$$\lim_{R \rightarrow \infty} V_{E_R}(z) = 0,$$

where we define $E_R = E \cap \{z \in \mathbb{C}: |z| \leq R\}$.

3. If sequences (P_n) of polynomials and $k_n \geq \deg(P_n)$ satisfy

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for all $z \in E$, then

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for all $z \in \mathbb{C}$.

4. $V_E(z) \equiv 0$.

In \mathbb{C}^m ($m > 1$), we cannot reduce the definition of non-thinness of E at ∞ to the non-thinness at 0 of some other sets E^* , as it was when $m = 1$. However, Proposition 1 suggests a way to define non-thinness at ∞ in higher dimensions:

Definition 1. Let E be a subset of \mathbb{C}^m . For any $R > 0$ define $E_R = E \cap \{z \in \mathbb{C}^m: |z| \leq R\}$. We say that E is non-thin at ∞ if

$$\lim_{R \rightarrow \infty} V_{E_R}(z) = 0$$

for all $z \in \mathbb{C}^m$.

If E is not non-thin at ∞ then we say E is thin at ∞ .

This definition of non-thinness at ∞ has good properties: If $E \subset \mathbb{C}^m$ is non-thin at ∞ and A is pluripolar then $E \setminus A$ is non-thin at ∞ ; if $E \subset \mathbb{C}^m$ and $F \subset \mathbb{C}^n$ are arbitrary sets, then E and F are non-thin at ∞ iff $E \times F \subset \mathbb{C}^m \times \mathbb{C}^n$ is non-thin at ∞ (see Lemma 2).

Our first result in this paper is the following, which is an analog to the case $m = 1$:

Theorem 2. *Let E be a closed subset of \mathbb{C}^m . Then the following two statements are equivalent:*

1. E is non-thin at ∞ .
2. If sequences (P_n) of polynomials and $k_n \geq \deg(P_n)$ satisfy

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for all $z \in E$, then

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for all $z \in \mathbb{C}^m$.

If E is non-thin at ∞ then it is easy to see that $V_E \equiv 0$. The converse is not true for $m \geq 2$, by the following example (which essentially is the same as Example 1.1 in [2]).

Example 1. Let E be a subset of \mathbb{C}^2 defined by

$$E = \{(z_1, z_2) \in \mathbb{C}^2: |z_2| \leq 1\} \cup \{(z_1, z_2) \in \mathbb{C}^2: z_1 = 0\}.$$

Then $V_E \equiv 0$ but E is thin at ∞ .

Proof. Using arguments in Example 1.1 in [2], it is easy to see that $V_E \equiv 0$.

If E was non-thin at ∞ then since $A = \{(z_1, z_2) \in \mathbb{C}^2: z_1 = 0\}$ is pluripolar, by Lemma 2 we should have $E \setminus A$ is non-thin at ∞ . In particular we should have $V_{E \setminus A} \equiv 0$. However as computed in [2] we have $V_{E \setminus A}(z) = \log^+ |z_2|$, which is a contradiction.

Hence E is thin at ∞ . \square

The following result gives a characterization of sets E with $V_E \equiv 0$, which also shows that the non-thin at ∞ sets are not rare.

Theorem 3. Let E be a closed subset of \mathbb{C}^m . Then $V_E \equiv 0$ iff any open neighborhood of E is non-thin at ∞ .

Example 2. Applying Theorem 3 to Example 1, we see that the set

$$\{(z_1, z_2) \in \mathbb{C}^2: |z_2| \leq 1\} \cup W,$$

is non-thin at ∞ in \mathbb{C}^2 , where W is an open neighborhood of the set $\{(z_1, z_2) \in \mathbb{C}^2: z_1 = 0\}$.

Example 3. Let Δ be a collection of complex lines L in \mathbb{C}^m such that

$$\bigcup_{L \in \Delta} L = \mathbb{C}^m.$$

Let E be a closed subset of \mathbb{C}^m such that for each $L \in \Delta$ the set $E \cap L$ considered as a subset in the one-dimensional complex line L is non-thin at ∞ . Then E is non-thin at ∞ as a subset in \mathbb{C}^m .

Proof. By Theorem 2 we need only to show that: If (P_n) is a sequence of polynomials, and $k_n \geq \deg(P_n)$ such that

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for $z \in E$, then

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for all $z \in \mathbb{C}^m$.

Let $w \in \mathbb{C}$ be a coordinate for L such that the coordinates z_1, \dots, z_n of \mathbb{C}^m are linear functions of w when restricted on L , and denote by $P_{n,L}(w)$ the restriction of P_n to L . Then $P_{n,L}(w)$ is a polynomial in one complex variable w of degree $\deg(P_{n,L}(w)) \leq \deg(P_n) \leq k_n$.

Then since

$$\limsup_{n \rightarrow \infty} |P_{n,L}(w)|^{1/k_n} \leq 1$$

for $w \in E \cap L$, and $E \cap L$ is non-thin at ∞ in the complex line L , hence

$$\limsup_{n \rightarrow \infty} |P_n(w)|^{1/k_n} \leq 1$$

for all $w \in L$. Since this is true for any complex line $L \in \Delta$, it is also true for their union, which is equal to \mathbb{C}^m . \square

In Theorem 2 of [3], we prove a Phragmen–Lindelof type inequality for subsets of \mathbb{C} having “big” capacity (by “big” capacity we mean a set E for which $C(E_R)$ grows proportionally to $\log R$). In the sequel, we state and prove a similar result for higher dimensions (see the remarks at the end of this section for more discussion about this).

Let $K \subset \mathbb{C}^m$ be compact. Then the Robin constant of K is defined as (see [6])

$$\gamma(K) = \limsup_{z \rightarrow \infty} [V_E^*(z) - \log |z|],$$

and the \mathbb{C}^m -capacity of K is

$$C(K) = e^{-\gamma(K)}. \quad (1.1)$$

Let $K \subset \{z \in \mathbb{C}^m: |z| \leq s\}$ be compact and non-pluripolar, where $s > 0$. Then for any $0 < t < 1$ we have

$$\sup_{|z| \leq s} V_K^*(z) \leq m \frac{1+t}{(1-t)^{2m-1}} \left(\log s + \log \frac{1}{t} - \log C(K) \right). \quad (1.2)$$

In fact, the proof of theorem in [8, p. 319] gives that

$$\int_{S^{2m-1}} V_K^*(rz) d\sigma(z) \leq m(\log r - \log C(K)),$$

for all $r \geq s$, where $d\sigma$ is the normalized area measure on the unit sphere S^{2m-1} in \mathbb{C}^m . Since V_K^* is a non-negative PSH function, if in the above inequality we choose $r = s/t$ and apply Harnack inequality (see for example Chapter 3 in [1]), we obtain

$$\sup_{|z| \leq s} V_K^*(z) \leq \frac{1+t}{(1-t)^{2m-1}} \int_{S^{2m-1}} V_K^*(sz/t) d\sigma(z) \leq m \frac{1+t}{(1-t)^{2m-1}} \left(\log s + \log \frac{1}{t} - \log C(K) \right).$$

Lemma 1. Let $E \subset \mathbb{C}^m$ be closed and satisfy the following condition: there exists $0 < \beta \leq 1$ such that

$$\limsup_{R \rightarrow \infty} \frac{\log C(E_R)}{\log R} \geq \beta > 1 - \frac{1}{m}, \quad (1.3)$$

where $C(E_R)$ is determined from formula (1.1).

Let (P_n) be a sequence of polynomials and $k_n \geq \deg(P_n)$. Assume that for all $z \in E$ we have

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq h(|z|)$$

where h is a positive real function such that

$$\limsup_{R \rightarrow \infty} \frac{\log h(R)}{\log R} \leq \tau < \infty. \quad (1.4)$$

For all $\lambda \in S^{2m-1}$, we define

$$\exp \left\{ \limsup_{n \rightarrow \infty} \frac{1}{2\pi k_n} \int_0^{2\pi} \log |P_n(e^{i\theta} \lambda)| d\theta \right\} =: C_\lambda.$$

Then for any $w \in \mathbb{C}$ and $\lambda \in S^{2m-1}$ we have

$$\limsup_{n \rightarrow \infty} |P_n(w\lambda)|^{1/k_n} \leq C_\lambda (1 + |w|)^{\tau/[1-m(1-\beta)]}.$$

Corollary 1. Let $E \subset \mathbb{C}^m$ be closed. Then E is non-thin at ∞ if it satisfies the following condition: There exists $0 < \beta \leq 1$ such that

$$\limsup_{R \rightarrow \infty} \frac{\log C(E_R)}{\log R} \geq \beta > 1 - \frac{1}{m},$$

where $C(E_R)$ is determined from formula (1.1).

Corollary 2. Let Δ be a non-empty closed subset of \mathbb{P}^{m-1} and $F = \bigcup_{\lambda \in \Delta} L_\lambda$ where L_λ is the complex line going through 0 with direction λ . Assume that

$$\limsup_{R \rightarrow \infty} \frac{\log C(F_R)}{\log R} = \gamma > 1 - \frac{1}{m},$$

where as before $F_R = F \cap \{z \in \mathbb{C}^m: |z| \leq R\}$.

Let $E \subset \mathbb{C}^m$ be closed and satisfy the following condition: there exists a constant $0 < \beta \leq 1$ with

$$m(1 - \gamma) < \beta$$

such that

$$\liminf_{s \rightarrow \infty} \frac{1}{\log s} \frac{1}{2\pi} \int_0^{2\pi} V_{E_s}^*(se^{it}\lambda) dt \leq 1 - \beta \quad (1.5)$$

for all $\lambda \in \Delta$. Then E is non-thin at ∞ .

Interesting questions may arise such as whether we have a Wiener criterion for non-thinness at ∞ in \mathbb{C}^m , or can we extend the above results to sequences of entire functions of bounded orders as done in [3] for one-dimensional. An example in [5], giving lower bounds for a constant in another result of Taylor in the cited paper [8], suggests that the best lower bound for β in Lemma 1 and Corollary 1 may be $1 - 1/m$. If the previous claim is true, it means that there are sets in \mathbb{C}^m with “big” capacities yet being thin at ∞ when $m > 1$. We hope to return to these issues in a future paper.

The rest of this paper is devoted to proving the results stated above.

2. Proofs of the results

Lemma 2.

- (a) Let E be a subset of \mathbb{C}^m . If E is non-thin at ∞ and A is pluripolar then $E \setminus A$ is non-thin at ∞ .
 (b) Let $E \subset \mathbb{C}^m$ and $F \subset \mathbb{C}^n$ be arbitrary sets. Then $E \times F \subset \mathbb{C}^m \times \mathbb{C}^n$ is non-thin at ∞ iff $E \subset \mathbb{C}^m$ and $F \subset \mathbb{C}^n$ are non-thin at ∞ .

Proof. (a) Define $F = E \setminus A$. For any $R > 0$ the set $E_R = E \cap \{z \in \mathbb{C}^m: |z| \leq R\}$ is bounded. Hence by Corollary 5.2.5 in [4] we have

$$V_{E_R}^* = V_{F_R}^*.$$

Fix a sequence $R_n \rightarrow \infty$, using the same argument as in the proof of Theorem 2 (see below), there exists a pluripolar set B such that

$$\lim_{n \rightarrow \infty} V_{F_{R_n}}(z) = \lim_{n \rightarrow \infty} V_{F_{R_n}}^*(z) = \lim_{n \rightarrow \infty} V_{E_{R_n}}^*(z) = \lim_{n \rightarrow \infty} V_{E_{R_n}}(z) = 0,$$

for all $z \in \mathbb{C}^m \setminus B$. Since $V_{F_{R_n}}(z)$ are PSH, the previous equality is then also true for $z \in B$, thus proves (a).

(b) Let $R > 0$. Then one can check easily that

$$\max\{V_{E_R}(z), V_{F_R}(w)\} \leq V_{E_R \times F_R}(z, w) \leq V_{E_R}(z) + V_{F_R}(w),$$

where $z \in \mathbb{C}^m$, $w \in \mathbb{C}^n$. This completes the proof of case (b). \square

Now we proceed to proving Theorem 2.

Proof of Theorem 2. In this proof fix a sequence $R_n \nearrow \infty$.

($2 \Rightarrow 1$) Since E is closed, for any $R > 0$ we have E_R is compact.

Fix $z_0 \in \mathbb{C}^m$. By Siciak's theorem (see Theorem 5.1.7 in [4]) there exists a sequence of polynomials (P_n) of degree (k_n) such that

$$\|P_n\|_{E_{R_n}}^{1/k_n} \leq 1,$$

for all $n = 1, 2, \dots$, and

$$\lim_{n \rightarrow \infty} \log |P_n(z_0)|^{1/k_n} = \lim_{n \rightarrow \infty} V_{E_{R_n}}(z_0). \quad (2.1)$$

Then for any $z \in E$ we have

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1.$$

Hence by assumption that Statement 2 is true, we have

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for all $z \in \mathbb{C}^m$. In particular, with $z = z_0$ we get from (2.1) that

$$\lim_{n \rightarrow \infty} V_{E_{R_n}}(z_0) \leq 0.$$

Since $z_0 \in \mathbb{C}^m$ is arbitrary and obviously $V_E(z) \geq 0$ for all z , we get Statement 1.

($1 \Rightarrow 2$) We use the ideas in [7]. Assume that Statement 1 is true. Consider any sequence (P_n) of polynomials and $k_n \geq \deg(P_n)$ such that

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1 \quad (2.2)$$

for all $z \in E$. For each n define

$$v_n(z) = \log |P_n(z)|^{1/k_n}.$$

Note that Statement 1 implies that for R large enough then E_R is non-pluripolar. Fix $R > 0$ such that E_R is non-pluripolar. For any $h, l \in \mathbb{N}$ define

$$E_R^{h,l} = \bigcap_{n=h}^{\infty} \left\{ z \in E_R : v_n(z) < \frac{1}{l} \right\}.$$

Then $E_R^{h,l} \subset E_R^{h+1,l}$ and from (2.2)

$$\bigcup_{h=1}^{\infty} E_R^{h,l} = E_R \quad (2.3)$$

for any $l \in \mathbb{N}$.

By definition of the pluricomplex Green function, for any $h, l \in \mathbb{N}$, $n \geq h$ and $z \in \mathbb{C}^m$

$$v_n(z) \leq V_{E_R^{h,l}}(z) + \frac{1}{l} \leq V_{E_R^{h,l}}^*(z) + \frac{1}{l}.$$

Hence taking the limsup of the above inequality as $n \rightarrow \infty$ we get

$$v(z) = \limsup_{n \rightarrow \infty} v_n(z) \leq V_{E_R^{h,l}}(z) + \frac{1}{l} \leq V_{E_R^{h,l}}^*(z) + \frac{1}{l},$$

for all $h, l \in \mathbb{N}$, $z \in \mathbb{C}^m$. Take the limit of this inequality as $h \rightarrow \infty$. Using (2.2), we see from Corollary 5.2.6 in [4] that

$$v(z) \leq \lim_{h \rightarrow \infty} V_{E_R^{h,l}}^*(z) + \frac{1}{l} = V_{E_R}^*(z) + \frac{1}{l},$$

for all $l \in \mathbb{N}$, $z \in \mathbb{C}^m$, $R > 0$. Since l is arbitrary, we get

$$v(z) \leq V_{E_R}^*(z), \quad (2.4)$$

for all $z \in \mathbb{C}^m$, $R > 0$.

For each $n \in \mathbb{N}$ define

$$A_n = \{ z \in \mathbb{C}^m : V_{E_{R_n}}(z) < V_{E_{R_n}}^*(z) \}$$

then A_n is pluripolar thus has Lebesgue measure zero, and $V_{E_{R_n}}(z) = V_{E_{R_n}}^*(z)$ for $z \in \mathbb{C}^m \setminus A_n$. Hence

$$A = \bigcup_{n=1}^{\infty} A_n$$

is also pluripolar and of Lebesgue measure zero, and we have $V_{E_{R_n}}(z) = V_{E_{R_n}}^*(z)$ for $z \in \mathbb{C}^m \setminus A$ and $n \in \mathbb{N}$. Hence for $z \in \mathbb{C}^m \setminus A$, applying (2.4) we get

$$v(z) \leq \lim_{n \rightarrow \infty} V_{E_{R_n}}^*(z) = \lim_{n \rightarrow \infty} V_{E_{R_n}}(z) = 0$$

for all $z \in \mathbb{C}^m \setminus A$. Since A is of Lebesgue measure zero, by definition of $v(z)$, we see that $v(z) \leq 0$ for all $z \in \mathbb{C}^m$. This completes the proof of Theorem 2. \square

Now we prove Theorem 3.

Proof of Theorem 3. (\Rightarrow) Let E be a closed subset of \mathbb{C}^m . Assume that $V_E \equiv 0$. We will show that any open neighborhood of E is non-thin at ∞ . Let F be any open neighborhood of E . Then for any $R > 0$ since E_R is compact, F is open and contains E_R , we can find $1 > \epsilon = \epsilon_R > 0$ such that

$$E_{R,\epsilon} = \{ z \in \mathbb{C}^m : \text{dist}(z, E_R) \leq \epsilon \} \subset F.$$

By Corollary 5.1.5 in [4] we have $V_{E_{R,\epsilon}}^*(z) = 0$ for $z \in E_{R,\epsilon} \supset E_R$. Now it is obvious that $E_{R,\epsilon} \subset F_{R+1}$ hence for $z \in E_R$

$$V_{F_{R+1}}^*(z) = 0. \quad (2.5)$$

Define

$$v(z) = \lim_{R \rightarrow \infty} V_{F_R}^*(z),$$

then v is the limit of a decreasing sequence of PSH functions hence v is itself PSH. By (2.5) for $z \in E$ we have $v(z) \equiv 0$. Thus by definition of the pluricomplex Green function

$$v(z) \leq V_E(z) = 0$$

for all $z \in \mathbb{C}^m$. This shows that F is non-thin at ∞ .

(\Leftarrow) Assume that E is an arbitrary subset of \mathbb{C}^m with $V_E \not\equiv 0$. Then $V_E(z_0) > 0$ for some $z_0 \in \mathbb{C}^m$, hence by definition of the pluricomplex Green function, there exists a function $u \in \mathbb{L}$ such that $u(z) \leq 0$ for $z \in E$, and $u(z_0) > 0$. Define

$$F = \{z \in \mathbb{C}^m : u(z) < u(z_0)/2\}.$$

F is open because u is upper-semicontinuous, and $E \subset F$ because $u|_E \leq 0$ and $u(z_0) > 0$. Now $u(z) < u(z_0)/2$ for $z \in F$, hence we have

$$u(z) \leq V_F(z) + u(z_0)/2$$

for all $z \in \mathbb{C}^m$. In particular choose $z = z_0$; we have $V_F(z_0) \geq u(z_0)/2 > 0$, hence F is thin at ∞ . \square

We conclude this section presenting the proofs of Lemma 1 and Corollaries 1 and 2.

Proof of Lemma 1. Define

$$u(z) = \limsup_{n \rightarrow \infty} \log |P_n(z)|^{1/k_n},$$

and let $u^*(z)$ be its upper-semicontinuous regularization. Then $u^*(z) \in \mathbb{L}$.

Now we define

$$\kappa_0 = \limsup_{z \in \mathbb{C}^m, z \rightarrow \infty} \frac{u^*(z)}{\log |z|}. \quad (2.6)$$

For any $R > 0$, by the definition of the pluricomplex Green function

$$u^*(z) \leq \kappa_0 V_{E_R}(z) + \log h(R). \quad (2.7)$$

For any $\lambda \in S^{2m-1}$ define

$$\kappa(\lambda) = \limsup_{w \in \mathbb{C}, w \rightarrow \infty} \frac{u(w\lambda)}{\log |w|} \leq \kappa_0.$$

Fix $\lambda \in S^{m-1}$. By definition

$$u(w\lambda) = \limsup_{n \rightarrow \infty} \frac{1}{k_n} \log |P_n(w\lambda)|$$

for all $w \in \mathbb{C}$. Hence using (2.7) and (1.2), for any $s > 0$, $0 < t < 1$ we have

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{k_n} \log |P_n(se^{i\theta}\lambda)| d\theta &\leq \frac{1}{2\pi} \int_0^{2\pi} u(se^{i\theta}\lambda) d\theta \\ &\leq \kappa_0 \frac{1}{2\pi} \int_0^{2\pi} V_{E_s}(se^{i\theta}\lambda) d\theta + \log h(s) \\ &\leq \kappa_0 m \frac{1+t}{(1-t)^{2m-1}} \left(\log s + \log \frac{1}{t} - \log C(E_s) \right) + \log h(s). \end{aligned}$$

By the previous inequality and (1.4), we get for any $0 < t < 1$:

$$\liminf_{s \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{1}{\log s} \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{k_n} \log |P_n(se^{i\theta}\lambda)| d\theta \leq \kappa_0 m \frac{1+t}{(1-t)^{2m-1}} (1-\beta) + \tau. \quad (2.8)$$

Taking the limit $t \rightarrow 0^+$ in the previous inequality we obtain

$$\liminf_{s \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{1}{\log s} \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{k_n} \log |P_n(se^{i\theta}\lambda)| d\theta \leq \kappa_0 m (1-\beta) + \tau. \quad (2.9)$$

Then by Lemma 2 in [3] we have

$$\kappa(\lambda) \leq \kappa_0 m (1-\beta) + \tau. \quad (2.10)$$

Since (2.10) is true for any $\lambda \in S^{2m-1}$, by definition (2.6) and Lemma 2 in [3] we have

$$\kappa_0 \leq \kappa_0 m(1 - \beta) + \tau,$$

or equivalently

$$\kappa_0 \leq \frac{\tau}{1 - m(1 - \beta)}.$$

From the above inequality, using Lemma 2 in [3] we get the conclusion of Lemma 1. \square

Proof of Corollary 1. Let (P_n) be a sequence of polynomials and let $k_n \geq \deg(P_n)$ satisfying

$$u(z) = \limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for all $z \in E$. Applying Lemma 1 for the set E , the sequences (P_n) and (k_n) , and $\tau = 0$, we get

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1,$$

for all $z \in \mathbb{C}^m$. Then Theorem 2 implies that E is non-thin at ∞ . \square

Proof of Corollary 2. From the assumptions, we have that E_R is non-pluripolar for $R > 0$ large enough.

Let (P_n) be a sequence of polynomials and let $k_n \geq \deg(P_n)$ satisfying

$$u(z) = \limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1$$

for all $z \in E$. Define

$$\kappa_0 = \limsup_{z \in \mathbb{C}^m, |z| \rightarrow \infty} \frac{u^*(z)}{\log |z|} \leq 1,$$

and

$$\kappa_1 = \sup_{\lambda \in \Delta} \limsup_{w \in \mathbb{C}, |w| \rightarrow \infty} \frac{u^*(w\lambda)}{\log |w|} \leq \kappa_0.$$

Applying Lemma 1 for the set F , the sequences (P_n) and (k_n) , and $\tau = \kappa_1$ we obtain

$$\kappa_0 \leq \frac{\kappa_1}{1 - m(1 - \gamma)}.$$

Using assumption (1.5), apply formula (2.10) for all $\lambda \in \Delta$, for the set E , the sequences (P_n) and (k_n) , and $\tau = 0$; we obtain

$$\kappa_1 \leq \kappa_0(1 - \beta).$$

Hence we have

$$\frac{\kappa_1}{1 - \beta} \leq \kappa_0 \leq \frac{\kappa_1}{1 - m(1 - \gamma)}.$$

From our assumption that $\beta > m(1 - \gamma)$, the above inequality implies that $\kappa_0 = \kappa_1 = 0$. Hence $u^*(z)$ is a constant. Moreover, $u^*(z) \leq 0$ on the set E which is non-pluripolar, hence $u^*(z) \leq 0$ for all $z \in \mathbb{C}^m$. By Theorem 2, E is non-thin at ∞ . \square

Acknowledgments

The author would like to thank Professor Eric Bedford and Professor Norman Levenberg for their generous and fruitful help. The author also would like to thank the referee for many helpful comments that helped to improve the paper.

References

- [1] Sheldon Axler, Paul Bourdon, Ramy Wade, Harmonic Function Theory, 2nd edition, Springer-Verlag, New York, 2001.
- [2] U. Cegrell, S. Kolodziej, N. Levenberg, Two problems on potential theory for unbounded sets, Math. Scand. 83 (1998) 265–276.
- [3] Dang Duc Trong, Tuyen Trung Truong, The growth at infinity of a sequence of entire functions of bounded orders, Complex Var. Elliptic Equ. 53 (8) (2008) 717–743.
- [4] Maciej Klimek, Pluripotential Theory, Courier International Ltd., Scotland, 1991.
- [5] S. Kolodziej, The logarithmic capacity in \mathbb{C}^n , Ann. Polon. Math. 48 (3) (1988) 253–267.
- [6] Norman Levenberg, Capacities in several complex variables, PhD thesis, The University of Michigan, 1984.
- [7] J. Muller, A. Yavrian, On polynomials sequences with restricted growth near infinity, Bull. London Math. Soc. 34 (2002) 189–199.
- [8] B.A. Taylor, An estimate for an extremal plurisubharmonic function \mathbb{C}^n , in: Seminar P. Lelong–P. Dolbeaut–H. Skoda, in: Lecture Notes in Math., vol. 1028, Springer, Berlin, 1983, pp. 318–328.